

## Package of Care to be provided according to type of Health facility

Minimum Package of Care - Primary Health Center	Preferred Package of Care - Primary Health Center - Comprehensive Sites (Secondary/Tertiary Hospital)	Optimal Package of Care - Comprehensive Sites (Secondary/Tertiary Hospital)
<p style="text-align: center;"><b>Basic Antenatal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Physical Exam (including wt, assessing gestational age, blood pressure)</li> <li>▪ Vaccinations</li> <li>▪ Infant feeding counseling</li> <li>▪ STI screening</li> <li>▪ Malaria prophylaxis (IPT)</li> </ul>	<p style="text-align: center;"><b>Basic Antenatal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Physical Exam (including wt, assessing gestational age, blood pressure)</li> <li>▪ Vaccinations</li> <li>▪ Infant feeding counseling</li> <li>▪ STI screening</li> <li>▪ Malaria prophylaxis (IPT and bednets)</li> <li>▪ Anemia screening</li> <li>▪ TB screening</li> </ul>	<p style="text-align: center;"><b>Basic Antenatal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Physical Exam (including wt, assessing gestational age, blood pressure)</li> <li>▪ Vaccinations</li> <li>▪ Infant feeding counseling</li> <li>▪ STI screening and treatment</li> <li>▪ Malaria prophylaxis (IPT and bednets)</li> <li>▪ Anemia screening</li> <li>▪ TB screening</li> <li>▪ OI diagnosis and management</li> <li>▪ Multivitamin supplementation</li> </ul>
<p style="text-align: center;"><b>Counseling and Testing services available within ANC services</b></p> <ul style="list-style-type: none"> <li>▪ CT integrated and directly provided within ANC via group pre-test counseling and individualized post test counseling</li> <li>▪ On-site point of service bleeding and testing (some sites will rely on lab technicians to conduct rapid HIV test)</li> </ul>	<p style="text-align: center;"><b>Counseling and Testing services available within ANC services</b></p> <ul style="list-style-type: none"> <li>▪ CT integrated and directly provided within ANC via group pre-test counseling and individualized post test counseling</li> <li>▪ On-site point of service bleeding and testing performed by ANC nurse or counselor</li> </ul>	<p style="text-align: center;"><b>Counseling and Testing services available within ANC services</b></p> <ul style="list-style-type: none"> <li>▪ CT integrated and directly provided within ANC via group pre-test counseling and individualized post test counseling</li> <li>▪ On-site point of service bleeding and testing performed by ANC nurse or counselor</li> </ul>
<p style="text-align: center;"><b>Partner &amp; Family Counseling and Testing services</b></p> <ul style="list-style-type: none"> <li>▪ Women provided with letter of invitation encouraging partners to get HIV tested</li> <li>▪ services for partner testing provided in VCT</li> </ul>	<p style="text-align: center;"><b>Partner &amp; Family Counseling and Testing services</b></p> <ul style="list-style-type: none"> <li>▪ Women provided with letter of invitation encouraging partners to get HIV tested</li> <li>▪ Services for partner testing provided within ANC and linked to PMTCT client</li> <li>▪ Partner testing register available</li> <li>▪ Couples counseling provided</li> </ul>	<p style="text-align: center;"><b>Partner &amp; Family Counseling and Testing services</b></p> <ul style="list-style-type: none"> <li>▪ Women provided with letter of invitation encouraging partners to get HIV tested</li> <li>▪ services for partner testing provided within ANC and linked to PMTCT client</li> <li>▪ Partner register available</li> <li>▪ Couples counseling provided</li> </ul>
<p style="text-align: center;"><b>Testing and Counseling services for PMTCT available within maternities/ Labor&amp; Delivery</b></p> <ul style="list-style-type: none"> <li>▪ Maternity may not be available</li> </ul>	<p style="text-align: center;"><b>Testing and Counseling services for PMTCT available within maternities/ Labor&amp; Delivery</b></p> <ul style="list-style-type: none"> <li>▪ On-site point of service testing</li> <li>▪ Requires all L&amp;D nurses to be trained in testing and counseling services</li> <li>▪ PMTCT Register in maternity setting that includes T&amp;C services</li> </ul>	<p style="text-align: center;"><b>Testing and Counseling services for PMTCT available within maternities/ Labor&amp; Delivery</b></p> <ul style="list-style-type: none"> <li>▪ On-site point of service testing</li> <li>▪ Requires all L&amp;D nurses to be trained in testing and counseling services</li> <li>▪ PMTCT Register in maternity setting that includes T&amp;C services</li> <li>▪ Retesting of negative pregnant women with test results &gt;3mos</li> </ul>
<p style="text-align: center;"><b><u>PMTCT Prophylaxis</u></b></p> <p><b>Mother:</b></p> <ul style="list-style-type: none"> <li>▪ Sd-NVP dispensed in ANC at first point of contact and documented in PMTCT register</li> <li>▪ Sd-NVP dispensed in L&amp;D ward for HIV positive</li> </ul>	<p style="text-align: center;"><b><u>PMTCT Prophylaxis</u></b></p> <p><b>Mother:</b></p> <ul style="list-style-type: none"> <li>▪ Referral to lab for CD4 blood draw or onsite blood draw in ANC for CD4 evaluation</li> <li>▪ Specimens sent to referral lab for processing</li> </ul>	<p style="text-align: center;"><b><u>PMTCT Prophylaxis</u></b></p> <p><b>Mother:</b></p> <ul style="list-style-type: none"> <li>▪ CD4 blood draw done on-site on day of HIV positive test result</li> <li>▪ Clinical Staging of all HIV positive pregnant women</li> </ul>

## Package of Care to be provided according to type of Health facility

Minimum Package of Care - Primary Health Center	Preferred Package of Care - Primary Health Center - Comprehensive Sites (Secondary/Tertiary Hospital)	Optimal Package of Care - Comprehensive Sites (Secondary/Tertiary Hospital)
<p>women</p> <ul style="list-style-type: none"> <li>▪ Consider 7 day post partum tail of AZT/3TC if available</li> </ul> <p><b>Exposed Infant:</b></p> <ul style="list-style-type: none"> <li>▪ Sd-NVP for HIV exposed infant</li> </ul>	<ul style="list-style-type: none"> <li>▪ AZT starting at 28weeks and sd-NVP provided to mom when she initiates her AZT regimen with instructions on how to self administer NVP during onset of labor (if anemic, d4T in place of AZT if available; alternatively sd-NVP plus 7 day post partum tail of AZT/3TC)</li> <li>▪ Referral to ART clinic for ART initiation if CD4&lt;350</li> <li>▪ Referral to ART clinic for pCTX for women with CD4&lt;250</li> </ul> <p><b>Exposed Infant:</b></p> <ul style="list-style-type: none"> <li>▪ Sd-NVP for HIV exposed infant plus AZT syrup for 7days – 4 weeks</li> </ul>	<ul style="list-style-type: none"> <li>▪ AZT starting at 28weeks and sd-NVP provided to mom when she initiates her AZT regimen with instructions on how to self administer NVP during onset of labor. (if anemic, d4T in place of AZT)</li> <li>▪ Post partum tail of AZT/3TC tail for 7days</li> <li>▪ ART for eligible pregnant and lactating women provided on-site</li> <li>▪ CTX Prophylaxis for women with CD4&lt;250</li> </ul> <p><b>Exposed Infant:</b></p> <ul style="list-style-type: none"> <li>▪ Sd-NVP for HIV exposed infant plus AZT syrup for 7days – 4 weeks</li> </ul>
<p><b>Mother Support Activities</b></p> <ul style="list-style-type: none"> <li>▪ Mother Support Groups</li> </ul>	<p><b>Mother Support Activities</b></p> <ul style="list-style-type: none"> <li>▪ Mother Support Groups</li> <li>▪ Infant feeding support groups (including hygiene)</li> </ul>	<p><b>Mother Support Activities</b></p> <ul style="list-style-type: none"> <li>▪ Mother Support Groups</li> <li>▪ Infant feeding support groups (including hygiene)</li> </ul>
<p><b>Post Natal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Identify/Treat puerperal infections</li> <li>▪ Vitamin A supplementation within one month pp (one dose 200,000 IU)</li> <li>▪ Referral to family planning services (counseling on family planning &amp; birth spacing)</li> <li>▪ Counseling on breast health and maternal nutrition</li> </ul>	<p><b>Post Natal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Identify/Treat puerperal infections</li> <li>▪ Vitamin A supplementation within one month pp (one dose 200,000 IU)</li> <li>▪ Family planning services linked to PMTCT services</li> <li>▪ Counseling on breast health and maternal nutrition</li> <li>▪ Counseling on CD4 re-screening for lactating women not on ART</li> </ul>	<p><b>Post Natal Care Services</b></p> <ul style="list-style-type: none"> <li>▪ Identify/Treat puerperal infections</li> <li>▪ Vitamin A supplementation within one month pp (one dose 200,000 IU)</li> <li>▪ Family planning services integrated into postnatal follow-up (counseling and provision of FP methods and birth spacing)</li> <li>▪ Counseling on breast health and maternal nutrition</li> <li>▪ Coordinated with infant follow-up visits</li> <li>▪ CD4 re-screening for lactating women not on ART</li> </ul>
<p><b>Follow-up of Exposed infants</b></p> <ul style="list-style-type: none"> <li>▪ pCTX at 6 weeks</li> <li>▪ Growth monitoring (weight)</li> <li>▪ Immunizations (provided by nurses or follow-up visits coordinated with immunization clinic)</li> <li>▪ Vitamin A supplementation at</li> <li>▪ Antibody testing at &gt;12 months or &gt;6wks post weaning</li> <li>▪ Infant feeding counseling</li> </ul>	<p><b>Follow-up of Exposed infants</b></p> <ul style="list-style-type: none"> <li>▪ pCTX at 6 weeks</li> <li>▪ Growth monitoring (weight, height, head circumference)</li> <li>▪ Immunizations provided by nurses</li> <li>▪ Vitamin A supplementation</li> <li>▪ Antibody testing at &gt;12 months or &gt;6wks post weaning</li> <li>▪ Referral for early infant diagnosis (DBS blood collection in some sites)</li> <li>▪ Infant feeding counseling</li> <li>▪ Insecticide treated bednets provided</li> <li>▪ Referral to care and treatment for HIV infected children</li> </ul>	<p><b>Follow-up of Exposed infants</b></p> <ul style="list-style-type: none"> <li>▪ pCTX at 6 weeks</li> <li>▪ Growth monitoring (weight, height, head circumference)</li> <li>▪ Immunizations provided by nurses</li> <li>▪ Vitamin A supplementation</li> <li>▪ Early infant diagnosis (DBS collected by nurses)</li> <li>▪ Antibody testing at &gt;12 months or &gt;6wks post weaning</li> <li>▪ Disease staging</li> <li>▪ Infant feeding counseling</li> <li>▪ Insecticide treated bednets provided</li> <li>▪ Care of infected children integrated into consultation</li> </ul>